AD					

AWARD NUMBER: W81XWH-12-1-0031

TITLE: Research in Prevention and Treatment of Noise-Induced Hearing Loss (NIHL)

PRINCIPAL

INVESTIGATOR: Kathleen Campbell, PhD

CONTRACTING

ORGANIZATION: Southern Illinois University School of Medicine

Springfield, IL 62794-9629

REPORT DATE: April 2014

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION

STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DO		Form Approved OMB No. 0704-0188				
data needed, and completing and reviewing this collection this burden to Department of Defense, Washington Heads	of information. Send comments regularters Services, Directorate for Infograny other provision of law, no perso	arding this burden estimate or an rmation Operations and Reports n shall be subject to any penalty	y other aspect of this o (0704-0188), 1215 Jeft	ching existing data sources, gathering and maintaining the ollection of information, including suggestions for reducing ferson Davis Highway, Suite 1204, Arlington, VA 22202- th a collection of information if it does not display a currently		
1. REPORT DATE	2. REPORT TYPE	NE33.	3.	DATES COVERED		
April 2014	Annual		11	April 2013 – 31 March 2014		
4. TITLE AND SUBTITLE	Timuu		CONTRACT NUMBER			
Research in Prevention and Trea	atment of Noise Indu	iced		GRANT NUMBER		
		81XWH-12-1-0031 PROGRAM ELEMENT NUMBER				
Hearing Loss (NIHL)						
6. AUTHOR(S)			5d	PROJECT NUMBER		
Kathleen Campbell, PhD	5e.	TASK NUMBER				
E-Mail: kcampbell@siumed.ed	5f.	WORK UNIT NUMBER				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)				PERFORMING ORGANIZATION REPORT		
Southern Illinois University Sch	ool of Medicine					
Springfield, IL 62794-9629						
a apanaapina (Manitapina Aasha	V NAME(O) AND ADDRES	0/50)	40	ODONOOD/MONITODIO AODONIVAVO		
9. SPONSORING / MONITORING AGENC U.S. Army Medical Research ar			10.	SPONSOR/MONITOR'S ACRONYM(S)		
Fort Detrick, Maryland 21702-		iu				
Tort Detrick, War yland 21/02-	7012		11.	11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STAT	FRACUT					
		4				
Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES						
	11:1 ::0 5	(D) 1 1	1.1.1.1.1	0 1		
				oprotective response from steady state		
and impulse noise-induced hearing loss (NIHL). The second year was dedicated to determining the latest time D-met may be						
administered for efficient protection. Groups of <i>Chinchillas laniger</i> (n = 10) received five intraperitoneal D-met (200 mg/kg/dose)						
every 12 hours beginning 7, 9, 12, 18, 34, 36, or 48 hours after steady state or impulse. D-met otoprotection was assessed by						
auditory brainstem response (ABR) analysis and outer hair cell (OHC) quantification. ABR analyses identified time-dependent D-						
met protection from steady state and impulse noise exposures when D-met administration began as late as 24 hours post-noise						
exposure. Histological cytocochleog	gram analysis is still in	progress to confirm ϵ	electrophysiolo	gical assessment. Statistical analysis		
is currently in progress.						
15. SUBJECT TERMS						
none provided						
		T .=	T			
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON		

a. REPORT

U

b. ABSTRACT

U

c. THIS PAGE

U

6

UU

USAMRMC

19b. TELEPHONE NUMBER (include area

Table of Contents

	<u>Page</u>
Introduction	4
Body	4
Key Research Accomplishments	5
Reportable Outcomes	5
Conclusion	5
References	N/A
Appendices	
Supporting Data	6

INTRODUCTION:

In military settings, noise exposure frequently exceeds the physical hearing protector capacity; resulting in Noise Induced Hearing Loss (NIHL). An oral antioxidant that could prevent permanent NIHL could allow more military personnel to be redeployed, to maintain the auditory capabilities needed in combat situations, and to keep our military personnel from having permanent disability. Currently, no pharmacologic protective agents exist in the clinical arena. We have developed an antioxidant, D-methionine (D-met), which has already shown promise in preventing NIHL in animals whether started before or after the noise exposure. In these studies we have performed dose timing studies in chinchillas to determine the latest time after noise exposure D-met may be administered to efficiently prevents impulse and steady state NIHL.

BODY:

Statement of Work (SOW): Year 2: Task: To determine maximum time delay for effective D-met post-noise rescue protection from permanent NIHL using auditory brainstem response (ABR) threshold shift and outer hair cell (OHC) measurements.

Methods:

Each animal group (9 groups for steady state noise and 9 groups for impulse noise) comprised 10 male *Chinchillas laniger*. To determine the latest protective D-met administration time post-noise, D-met started at time delays of 7, 9, 12, 18, 24, 36, 48, and 72 hours after completion of the steady state or impulse noise exposure and then another 48 hours BID. The control group received the placebo starting 7 hours after cessation of the noise. ABRs were measured at baseline, prior to any drug or noise exposure, and again 21 days after noise exposure cessation using tone-burst stimuli centered at the frequencies of 2, 4, 6, 8, 14 and 20 kHz. Threshold was defined as the lowest intensity capable of eliciting a replicable, visually detectable response.

Steady state noise exposure comprised a 105 dB SPL narrow band of noise centered at 4 kHz for 6 hours. Impulse noise consisted of simulated M-16 weapon fire at 155 dB peSPL 150 impulses at 2/s.

After the 21 day post-noise ABR assessment, each animal was sacrificed by decapitation and the cochleae harvested for OHC. Animals completing experimentation will be decapitated and cochleae perfused with fixative through the perilymphatic spaces. The primary fixative will be 10% formalin at 4 degrees C in PBS. A small hole was first hand-drilled into the otic capsule with a sharpened pick. *In vitro* perfusion was performed intermittently within 5 minutes of sacrifice through the small hole in the scala tympani, allowing the fluid to exit through the opened oval window. After perfusion fixation, the round window membrane was removed and the cochleae immersed in 10% formalin and placed in the refrigerator.

After cochlear preservation, the cochlear epithelium was carefully dissected in one half turn sections, stained with either hematoxylin (Eosin Y counterstain) or fluorescent markers (Alexafluor-488 Phalloidin Green), and mounted onto slides for cytocochleogram analysis using Cytogram software (Kresge Hearing Research Institute, University of Michigan; Ann Arbor).

Results:

Year 2 Steady State and Impulse NIHL Studies:

One hundred sixty of one hundred eighty of the scheduled animals were successfully tested for D-met protection from steady state and impulse NIHL. We determined no D-met protection with a 48 hour rescue dosing regimen and therefore did not test the 72 hour time point to prevent animal pain, suffering, and sacrifice.

Current steady state and impulse noise-induced threshold shift results at 21 day post-noise exposure (Figures 1 and 2, respectively) identifies D-met protection when rescue dosing was administered as late as 24 hours post-

noise exposure. D-met provided protection when it was administered between 7 and 24 hours after noise exposure. At 36 hours, threshold shifts from D-met-treated animals were still noticeably still less than control thresholds, but with little or no significance. Finally, the 48 hour rescue regimen measured very little protection compared to saline-injected controls.

Year 2 studies performed almost twice as many animal experiments than the previous year. Thus, statistical and histological analyses are pending and in progress. We will finalize Year 2 data while Year 3 experimentation planning and protocol approvals are developed and scheduled.

KEY RESEARCH ACCOMPLISHMENTS:

- 80/90 of the animals scheduled for testing D-met time-delayed protection from impulse NIHL were tested using saline or 200 mg/kg/dose D-met.
- 80/90 of the animals scheduled for testing D-met time-delayed protection from steady state NIHL were tested with either saline or 200 mg/kg/dose D-met.
- ABR analysis identifies time-delayed D-met protection from steady state and impulse NIHL when D-met administration begins as late as 24 hours post-noise exposure.
- Learned and developed a technique for cytocochleagram quantitative analysis of chinchilla cochlear hair cells.
- Developed an animal model for chinchilla cytocochleagram quantitative analysis.
- Dissected, stained and prepared slides for 40% of the specimens.
- Cochlear whole mounts and cytocochleogram analyses are currently under development and in progress.

REPORTABLE OUTCOMES:

We have extended our published 7-hour D-met rescue from steady state noise exposure (Campbell et al. 2011) to protection from steady state and impulse NIHL when D-met administration begins at least 24 hours postnoise exposure. The results have just recently been analyzed, require statistical analysis, and have not yet been presented or published.

CONCLUSION:

We have identified a 24 hour time-delayed protective D-met dose from steady state and impulse NIHL. The 36 hour dosing regimen continues to yield threshold shifts lower than controls, but they are also greater than the 24-hour dosing group. Thus, the grant's third year will design animal experiments that test extended time-delayed D-met administration beginning between 24 and 36 hours, possibly at 28 and 32 hours, to refine the latest possible time D-met administration may begin to protect from steady state or impulse NIHL. These data will enable us to proceed to FDA approved clinical trials submit the data for publication and to the FDA.

REFERENCES:

Campbell, K., Claussen, A., Verhulst, S., Fox, D., Hughes, L. 2011. D-methionine (D-met) significantly rescues noise-induced hearing loss: timing studies. Hear Res. Dec: 282(1-2): 138-44.

APPENDICES:

N/A

SUPPORTING DATA:




Figure 1. Steady state noise-induced ABR threshold shifts in chinchillas 21 days after a 105dB peSPL 6 hour noise exposure centered at 4 kHz (n = 10 per group). D-met has provided protection when administration began as late as 24 hours post-noise exposure. Statistical analysis is currently in progress.

Time-Delayed D-methionine (D-met) Rescue from Impulse Noise Exposure: ABR Threshold Shifts




Figure 2. Impulse noise-induced ABR threshold shifts in chinchillas 21 days after simulated M-16 weapon fire at 155 dB peSPL 150 impulses at 2/s (n = 10 per group). D-met has provided protection when administration began as late as 24 hours post-noise exposure. Statistical analysis is currently in progress.